

# Nitrogen retranslocation, allocation, and utilization in bare root *Larix olgensis* seedlings

WEI Hong-xu • XU Cheng-yang • MA Lü-yi • DUAN Jie  
JIANG Li-ni • REN Jun

Received: 2011-02-28; Accepted: 2011-04-13  
© Northeast Forestry University and Springer-Verlag Berlin Heidelberg 2012

**Abstract:** We quantified biomass accumulation and nitrogen (N) retranslocation, allocation, and utilization of Changbai larch (*Larix olgensis*) seedlings subjected to four fertilization treatments (24, 59, 81, 117 kg·ha<sup>-1</sup> N) with an unfertilized control during summer and autumn 2009. Ammonium phosphate (18-46-0) was the fertilizer used in all treatments. On both sampling dates, the needles had greater biomass and N content than new (2009) stems and old (2008) stems, and coarse, medium and fine roots (diameters of >5, 2–5 mm, and 0–2 mm, respectively). Higher N concentration was observed in old stems and coarse roots than that in new stems and medium roots. In mid-summer, fine roots had higher N concentration than coarse roots. The treatment with 24 kg·ha<sup>-1</sup> N had the greatest biomass and N content in needles and old stems, and highest net N retranslocation (NRA) and amount of N derived from soil. On September 21, no N translocation was observed, while the treatment with 24 kg·ha<sup>-1</sup> N had the highest N utilization efficiency and fertilizer efficiency. Vector analysis revealed that all four fertilization treatments induced N

excess relative to the control. The treatments with 59, 81, 117 kg·ha<sup>-1</sup> N induce N excess compared with treatments at 24 kg·ha<sup>-1</sup> N. We conclude that the traditional local fertilizer application rates exceeded N requirements and N uptake ability for Changbai larch seedlings. The application rate of 24 kg·ha<sup>-1</sup> N is recommended.

**Keywords:** *Larix olgensis*; Nitrogen; retranslocation; allocation; biomass; Vector analysis

## Introduction

Nitrogen (N) availability is one of the important factors influencing growth of tree seedlings (Salifu and Timmer 2001, 2003; Salifu et al. 2008, 2009; Bown et al. 2010). Storage of N is important for young trees because buds break in the spring is sub-optimal when N uptake by the roots before photosynthesis can provide carbon skeletons for amino acid synthesis (Millard and Nielsen 1989). Deciduous trees store N mainly in woody stems and roots, while evergreen trees store N mainly in shoots (Millard and Grelet 2010).

Retranslocation of stored nutrient is important for the production of new tissues at all stages of development from the seedling to the mature tree (Nambiar and Fife 1991). Studies have focused on nutrient retranslocation in juvenile evergreen conifer seedlings (Nambiar and Fife 1987, 1991; van den Driessche 1984; Imo and Timmer 2001; Salifu and Timmer 2001, 2003), hardwood seedlings (Salifu et al. 2008, 2009), and mature tree stands (Helmisaari 1992; Mead and Preston 1994; Chapin and Kedrowski 1983). The main conclusions from these reports include: (1) the leaf is the main source and sink of retranslocated nutrients in evergreen conifers (Nambiar and Fife 1987, 1991). (2) net retranslocation is mainly driven by the amount of plant N reserves established before planting (Nambiar and Fife 1987; Imo and Timmer 2001; Salifu and Timmer 2001, 2003). (3) retranslocation is more efficient in trees growing on less fertile soils (Miller et al. 1979; Imo and Timmer 2001; Salifu and Timmer 2001). It is, however, not clear whether nutrient retrans-

Foundation project: The study was financially supported by the Special Industry Project of Ministry of Finance (Grant No. 201104051) and “11<sup>th</sup> Five-Year Plant” National Key Technology and R&D Program from Chinese State Forestry Administration (Grant No. 2006BAD24B01).

The online version is available at <http://www.springerlink.com>

WEI Hong-xu • XU Cheng-yang (✉) • MA Lü-yi  
DUAN Jie • JIANG Li-ni

Key Laboratory of Silviculture and Conservation, Ministry of Education, Beijing Forestry University, Beijing 100083, China.

E-mail: [cyxu@bjfu.edu.cn](mailto:cyxu@bjfu.edu.cn)

WEI Hong-xu  
Flower R&D Center, Zhejiang Academy of Agricultural Sciences, Xiaoshan District, Hangzhou 311202, China.

Jun Ren  
Jilin Provincial Academy of Forestry Sciences, Changchun 133033, China.

Responsible editor: Yu Lei

location from nutrient reserves in the whole seedling, such as from combined stems and roots (CSR) in deciduous conifers, follows the same pattern. Most previous studies focused on N retranslocated from shoots only (Salifu and Timmer 2001).

In response to external resource changes, trees tend to allocate internal resources to the tissues responsible for acquiring the most limiting compounds (Welker et al. 1991; Lehtilä et al. 2000; Ueda et al. 2009). N allocation to shoots improves carbon acquisition for carbohydrate production, and photosynthetic capacity is strongly correlated with the N content of leaves (Evans 1989). The decrease in N concentration in green leaves may relate to an increase in the proportion of plant N invested in roots (Lutze and Gifford 1998). Mean N concentration was observed to be significantly higher in smaller roots (< 2 mm diameter) than in larger roots (2–5 mm) (Gordon and Jackson 2000). However, most research has focused on evergreen plants or plants in leaf. There has been little research on patterns of N allocation over seasons within deciduous woody stock.

Changbai larch (*Larix olgensis* H.) is an important commercial timber tree species in north-eastern China (Zhu et al. 2008). Prior to this study, the local fertilizer regime for growing Changbai larch seedlings appeared to result in sub-optimal seedling quality (unpublished data). Therefore, we investigated the growth and N translocation, allocation, and utilization of transplanted bare root Changbai larch seedlings to find an optimal fertilizer regime for this species. We hypothesized that Changbai larch seedlings would support growth of new tissues by N retranslocation from CSR rather than by N uptake from soils, and N would be allocated mostly to needles and fine roots.

## Materials and methods

Seeds of Changbai larch were collected by employees of Xiaobeihu Forest Station (128°28' E, 44°03' N) from a local wild stand in Heilongjiang province, northeast China. After the collection, the seeds were transported to Jiangmifeng nursery (43°45' N, 126°45' E), Jilin City, Jilin Province, north-eastern China, and stored at 0–4°C. In Jilin, annual precipitation is 650–750 mm, with less than 200 mm from May to mid-June. The average annual temperature is 3–5°C, and the average temperature in the early growing season ranges from 4 to 9°C.

Seeds were soaked in 5% potassium permanganate (W/W) solution for 24 h and stratified for 5 d at 0–4°C. Then the seeds were sowed in a nursery bed at the density of 700 seed·m<sup>-2</sup> in late April 2008. In mid June 2008, the germinated seedlings were thinned to about 550 seedling·m<sup>-2</sup>. After excavation from the seedling bed in October 2008 and cold storage until mid April 2009, the seedlings were transplanted into another nursery bed with initial properties of surface soil (0–20cm) determined (Bao 2000): pH of 6.08, electrical conductivity (25°C) of 0.15 dS·m<sup>-1</sup>, total N of 2.07 mg·g<sup>-1</sup>, NH<sub>4</sub><sup>+</sup>-N of 1.88 mg·kg<sup>-1</sup>, NO<sub>3</sub><sup>-</sup>-N of 277.98 mg·kg<sup>-1</sup> and organic C of 24.28 mg·g<sup>-1</sup>.

Locally prescribed fertilizer applications in 2008 included: ammonium phosphate (18-46-0) incorporated into surface soil before sowing at 54 kg·ha<sup>-1</sup> N in late April, and two top-dress

applications of a 1:1 combination of granular ammonium nitrate (21-0-0) and urea (46-0-0) at 65 and 75 kg·ha<sup>-1</sup> N in late June and early July, respectively. The experiment began in the spring of 2009. There were four fertilizer treatments (24, 59, 81, 117 kg·N·ha<sup>-1</sup>) and an unfertilized control treatment. Fertilizer was applied as ammonium phosphate (18-46-0), incorporated into surface soils in mid April, 2009. A fertilizer rate of 81 kg·N·ha<sup>-1</sup> is operationally employed for local seedling production. For each treatment, there were three randomly assigned plots (replicates). Each plot had an area of 1 m × 1 m with 0.1–0.2 m buffer between plots and plastic barriers inserted to a depth of 0.2 m within each buffer to eliminate lateral movement of fertilizer between plots. All plots shared the same irrigation regime performed by automatic sprayers installed along the seed beds. Weed control was conducted manually. Frequency and intensity of irrigation and weed control were adjusted based on seedling growth rates and weather conditions throughout the growing season.

On both July 2 and September 21, 2009, 150 seedlings (10 seedling·plot<sup>-1</sup> × 3 plot·treatment<sup>-1</sup> × 5 treatments) were randomly excavated from the nursery bed. All seedlings were divided into shoot and root fractions at the cotyledon scar. Shoots were divided into needles, old (2008) stems (dark brown), new (2009) stems (light brown). Roots were divided into coarse roots (>5 mm in diameter), medium roots (2–5 mm in diameter) and fine roots (0–2 mm in diameter). The color of coarse and medium roots was dark brown, and the color of most fine roots was white. For each tissue fraction, the ten samples per plot and treatment were bulked and then transported to the laboratory on ice (0–2°C), where roots were washed free of soil. Dry biomass was measured for each bulked tissue after oven-drying for 48 h at 70°C. Bulked tissue samples were ground to pass a 1 mm sieve, then one 0.2 g-sample for each tissue was digested in 5 mL of H<sub>2</sub>O<sub>2</sub>-H<sub>2</sub>SO<sub>4</sub> and diluted to 50 mL. Total seedling N concentration was measured with 5mL of digestion solution using an automatic N analyzer (UDK 152 automatic N analyzer, VELP Co., Usmate (MB), Italy).

Net N retranslocation amount (*NRA*) was computed as (Salifu and Timmer 2001):

$$NRA = A - B \quad (1)$$

Where, *A* is the total N content (storage) of the whole seedling before transplanting in mid April 2009 (old stem + intact root). In this study, *A*=14.51 mg·seedling<sup>-1</sup> N. *B* is the total N content in old tissues (old stem + dark brown root > 2 mm diameter) on July 2 or September 21. Following Salifu and Timmer (2001), the N content in dead tissue and N leached from seedlings were assumed negligible.

Net N retranslocation ratio (*NRR*) was computed as:

$$NRR = (A - B) \times A^{-1} \times 100\% \quad (2)$$

Net amount of N derived from soil (*ANDFS*) was computed as:

$$ANDFS = C - NRA \quad (3)$$

Where,  $C$  is the total N content in new tissues (needles + new stem + white fine roots in 0–2 mm diameter) on July 2 or September 21.

Net ratio of N derived from soil ( $RNDFS$ ) was computed as:

$$RNDFS = ANDFS \times D^{-1} \times 100\% \quad (4)$$

Where,  $D$  is the total N content in initial soil and fertilizer.

N utilization efficiency ( $U$ ) was computed as:

$$U = Bio \times D^{-1} \times 100\% \quad (5)$$

Where,  $Bio$  was the dry biomass of the whole seedling or combined stem and root on July 2 or September 21.

Fertilizer efficiency ( $FN$ ) was computed as:

$$FN = Bio \times E^{-1} \quad (6)$$

Where,  $E$  was total N content in fertilizer.

Seedling biomass, N concentration, and N content (biomass multiplied by N concentration) were averaged for the 10 seedlings per plot on each sampling date. These values, as well as  $NRR$ ,  $ANDFS$ ,  $RNDFS$ ,  $U$ , and  $FN$ , were analyzed using analysis of variance (ANOVA) based on the General Linear Model (GLM) procedure of SAS (9.0) (SAS Institute Inc., NC, USA). For analyses of variance, the effects of treatment ( $F$ ) and tissue ( $T$ ) (needles, old stems, new stems, and coarse, medium and fine

roots) were considered fixed. When treatment effects were significant, means were ranked according to Tukey's studentized range test at  $\alpha = 0.05$ . When the effects of  $T$  or  $F$  were significant, a one-way analysis of variance (ANOVA) was employed to compare means among tissue fractions or treatments, respectively. For biomass and N status data, the interaction of  $T$  and  $F$  was significant and a two-way ANOVA was employed. Instantaneous vector diagnosis (Salifu and Timmer 2001) was employed to facilitate interpretation of the effects of fertilization treatments on whole seedlings or on combined stem and root growth. Nutritional interpretations of directional vector changes in biomass, N content, and N concentration on September 21 follow the conventions of Salifu and Timmer (2001).

## Results

### Biomass, N concentration and N content

Fertilization significantly affected biomass and N content on July 2 (Table 1). Seedlings fertilized with  $24 \text{ kg} \cdot \text{ha}^{-1}$  N had the greatest biomass ( $0.58 \pm 0.14 \text{ g}$  and highest N content ( $11.89 \pm 4.08 \text{ mg}$ ), followed by seedlings fertilized with  $59 \text{ kg} \cdot \text{ha}^{-1}$  N ( $0.50 \pm 0.12 \text{ g}$  biomass and  $10.48 \pm 4.08 \text{ mg}$  N content). Seedlings fertilized with  $117 \text{ kg} \cdot \text{ha}^{-1}$  N and the control had the least biomass ( $0.44 \pm 0.12 \text{ g}$ ) and lowest N content ( $8.93 \pm 3.28 \text{ mg}$ ). On September 21, seedlings fertilized with  $24 \text{ kg} \cdot \text{ha}^{-1}$  N had the greatest biomass ( $1.61 \pm 0.26 \text{ g}$ ), followed by seedlings in the control treatment ( $1.50 \pm 0.20 \text{ g}$ ). Fertilization rate did not affect the ratio (R/S) of root and shoot biomass on either July 2 ( $p = 0.4964$ ) or September 21 ( $p = 0.3836$ ). R/S ranged between 0.22 and 0.26 in July, and 0.32 to 0.48 in September.

**Table 1.** ANOVA for seedling tissue ( $T$ ) and fertilization treatment ( $F$ ), and their interaction on biomass, and N concentration, and N content on July 2 and September 21, 2009.

| Date    | Source of variation | $df^*$ | Biomass             |          | N concentration |          | N content |          |
|---------|---------------------|--------|---------------------|----------|-----------------|----------|-----------|----------|
|         |                     |        | MS <sup>&amp;</sup> | $Pr > F$ | MS              | $Pr > F$ | MS        | $Pr > F$ |
| Jul. 2  | $T$                 | 5      | 4.76                | <0.0001  | 934.76          | <0.0001  | 3975.35   | <0.0001  |
|         | $F$                 | 4      | 0.05                | <0.0001  | 12.85           | 0.1728   | 26.47     | 0.0003   |
|         | $T \times F$        | 20     | 0.01                | <0.0001  | 8.67            | 0.3598   | 13.24     | 0.0003   |
|         | Error               | 60     | 0.004               | -        | 7.77            | -        | 4.23      | -        |
| Sep. 21 | $T$                 | 5      | 8.67                | <0.0001  | 279.12          | <0.0001  | 1276.42   | <0.0001  |
|         | $F$                 | 4      | 1.05                | 0.0008   | 0.57            | 0.5713   | 275.90    | 0.0572   |
|         | $T \times F$        | 20     | 0.22                | 0.3367   | 0.18            | 0.1784   | 137.44    | 0.2293   |
|         | Error               | 60     | 0.19                | -        | 36.93           | -        | 107.55    | -        |

\*  $df$ , degree of freedom. <sup>&</sup> MS, mean square.

On both sampling dates, biomass, N concentration, and N content differed significantly between tissues (Table 1). Needles had the greatest biomass and N content (Table 2). Compared with new stems, old stems had substantially more biomass and higher N content and N concentration. On July 2, the coarse roots (>5 mm in diameter) had most biomass, while the fine

roots (0–2 mm in diameter) had the highest N concentration and content. On September 21, fine roots had more biomass than coarse roots (2–5 mm diameter). Neither N concentration nor N content differed among three root diameter classes in September (Table 2).

The interaction of the effects of fertilization treatment and tis-


sue were significant for biomass and N content on July 2 only (Table 1). Needles had greatest biomass and highest N content in treatments fertilized with 24 kg·ha<sup>-1</sup> and 59 kg·ha<sup>-1</sup> N (Fig. 1A, C).

Old (2008) stem biomass was greatest in control (0) and 24 kg·ha<sup>-1</sup> N samples (Fig. 1B), while N content in old stems was highest at 24 kg·ha<sup>-1</sup> N (Fig. 1D).

**Table 2. Biomass and N concentration and content in seedling needles, new stems, old stems, and roots of diameters >5, 2–5, and 0–2 mm on July 2<sup>nd</sup> and September 21<sup>st</sup>, 2010. (mean ± SE)**

| Tissue     | Jul. 2      |                                       |                | Sep. 21     |                                       |                |
|------------|-------------|---------------------------------------|----------------|-------------|---------------------------------------|----------------|
|            | Biomass (g) | N concentration (mg·g <sup>-1</sup> ) | N content (mg) | Biomass (g) | N concentration (mg·g <sup>-1</sup> ) | N content (mg) |
| Needles    | 1.60±0.05a* | 26.88±0.58a                           | 42.89±1.54a    | 2.25±0.11a  | 13.52±0.63ab                          | 30.46±2.08a    |
| New stem   | 0.25±0.01b  | 12.54±1.10c                           | 6.63±0.59b     | 1.41±0.13b  | 5.72±0.41c                            | 8.08±0.91c     |
| Old stem   | 0.55±0.02c  | 18.31±0.96b                           | 4.68±0.39bc    | 2.18±0.20a  | 9.10±2.01bc                           | 20.89±5.49ab   |
| Root (>5)  | 0.05±0.01d  | 6.85±0.53d                            | 0.32±0.06d     | 0.73±0.07cd | 18.03±2.59a                           | 13.01±2.26bc   |
| Root (2–5) | 0.25±0.01c  | 5.41±0.32d                            | 1.33±0.06d     | 0.36±0.03d  | 14.51±1.47ab                          | 5.14±0.67c     |
| Root (0–2) | 0.27±0.01c  | 14.45±0.65c                           | 4.09±0.31c     | 1.19±0.15bc | 11.52±1.55abc                         | 13.32±2.86bc   |

\* Different letters in the same column indicate statistically significant differences according to Tukey's studentized range test at the 0.05 level.



**Fig. 1 Biomass (top) and N content (bottom) in needles (left) and old stems (right) of seedlings in different fertilization treatments on July 2, 2009.** Different letters above the bars indicate statistical differences according to Tukey's studentized range test at  $\alpha = 0.05$ .

#### N retranslocation

Fertilization significantly affected N retranslocation on July 2, but not September 21 (Table 3). On July 2, net N retranslocation amount (*NRA*) and ratio (*NRR*) were lowest in the 24 kg·ha<sup>-1</sup> N treatment, and the amount and ratio of N derived from soil (*ANDFS* and *RNDFS*, respectively) were highest in this treat-

ment (Table 3). On September 21, *NRA* and *NRR* values were negative, and *RNDFS* values were greater than 100%, indicating N gain rather than loss from the year-old tissues (Table 3).

#### N utilization

N utilization efficiencies of both CSR and whole seedlings were lowest at fertilization rates of 59 kg·ha<sup>-1</sup> N and 117 kg·ha<sup>-1</sup> N


(Fig. 2A, B). Among four fertilization treatments, fertilizer efficiencies for both CSR and whole seedlings were highest in the

24 kg·ha<sup>-1</sup> N treatment (Fig. 2 C,D).

**Table 3.** Net N retranslocation amount (*NRA*) and ratio (*NRR*) and amount of N derived from soil (*ANDFS*) and ratio of N derived from soil (*RNDFS*) in five treatments on sampling dates of July 2 and September 21, 2009. (mean ± SE)

| Investigating Date | Fertilizer treatment (kg·N·ha <sup>-1</sup> ) | <i>NRA</i> (mg·seedling <sup>-1</sup> ) | <i>NRR</i> (%)  | <i>ANDFS</i> (mg·seedling <sup>-1</sup> ) | <i>RNDFS</i> (%) |
|--------------------|---|---|-----------------|---|------------------|
| Jul. 2             | 0   | 8.38±0.09a <sup>a</sup>                 | 57.73±0.60a     | 39.07±4.80b                               | 81.99±1.72b      |
|                    | 24  | 2.64±0.34b                              | 18.21±2.32b     | 56.84±1.37a                               | 95.58±0.46a      |
|                    | 59  | 7.29±1.07a                              | 50.26±7.38a     | 48.44±2.63ab                              | 86.80±2.24b      |
|                    | 81  | 7.38±1.31a                              | 50.86±9.03a     | 41.47±2.54b                               | 85.15±1.61b      |
|                    | 117   | 5.43±0.66ab                             | 37.41±4.58ab    | 41.36±2.25b                               | 88.39±1.31ab     |
|                    | <i>Pr&gt;F</i>                                |   | 0.0046          | 0.0093                                    | 0.0014           |
| Sep. 21            | 0   | -81.50±18.71a                           | -561.70±128.95a | 170.90±43.44a                             | 194.14±8.65a     |
|                    | 24  | -86.59±24.51a                           | -596.77±168.89a | 161.94±39.36a                             | 212.41±15.20a    |
|                    | 59  | -47.54±13.70a                           | -327.66±94.39a  | 100.92±20.01a                             | 185.49±17.35a    |
|                    | 81  | -50.51±11.25a                           | -348.10±77.52a  | 118.67±22.25a                             | 172.38±5.49a     |
|                    | 117   | -33.27±1.83a                            | -229.27±12.59a  | 70.33±2.96a                               | 194.00±17.76a    |
|                    | <i>Pr&gt;F</i>                                |   | 0.1558          | 0.1646                                    | 0.4008           |

<sup>a</sup> Different letters in the same column indicate statistically differences according to Tukey's studentized range test at  $\alpha = 0.05$ .




**Fig. 2** N utilization efficiency (top) (biomass accumulation (g) /sum of available N content in initial soil and N fertilization amount (kg)) and fertilizer efficiency (bottom) (biomass accumulation (g) /total N content in fertilizer (kg)) in CSR (left) and whole seedlings (right) in different fertilization treatments on September 21, 2009. Different letters above the bars indicate statistical differences according to Tukey's studentized range test at  $\alpha = 0.05$ .

Vector analysis

On September 21 fertilization induced an N excess in both nee-

dles and CSR relative to the controls (Fig.3). Comparing the treatments with the control, the 81kg·ha<sup>-1</sup> and 117 kg·ha<sup>-1</sup> N resulted in N depletion in needles (Fig. 3A) and CSR (Fig. 3B). Compared with the 24 kg·ha<sup>-1</sup> N, all higher fertilization rates

induced N excess in both needles (Fig. 3A) and CSR (Fig. 3B).



**Fig. 3** Vector analysis of N concentration, N content, and biomass of seedlings sampled on September 21, 2009 in four fertilization treatments (24, 59, 81, and 117 kg·ha<sup>-1</sup> N) plus an unfertilized control (0N) in needles (A) and CSR (combined stems and roots) (B). Seedlings receiving no fertilizer (0N) or 24 kg·ha<sup>-1</sup> N (24N) were normalized to 100 compared to all fertilization treatments or higher rate fertilization treatments, respectively. Shift E and F indicate nutrient excess. Shift G indicates nutrient depletion (adapted from Salifu and Timmer 2003).

## Discussion

Negative effects on biomass of fertilization treatments at rates higher than 24 kg·ha<sup>-1</sup> N on both sampling dates may relate to the high initial NO<sub>3</sub><sup>-</sup>-N content in our soils (277.98 mg·kg<sup>-1</sup>). Levels of NO<sub>3</sub><sup>-</sup>-N in the Jianmifeng Nursery are high compared to those of other bare root seedling nurseries, for example 0.9 mg·kg<sup>-1</sup> (Islam et al. 2009). More biomass in needles of our Changbai larch (Table 2) concurred with carbon allocation pattern in saplings of another *Larix* species (Kagawa et al. 2006). More N allocation to needles than other tissues (Table 2) indicates a high photosynthetic capacity in Changbai larch. More biomass and N accumulation in old, woody stems than in new stems (Table 2) suggested an accumulation of N reserves in CSR as observed in other deciduous tree species (Weinbaum and van Kessel 1998; Millard et al. 1998, 2001; Dyckmans and Flessa 2001; Frak et al. 2002; Silla and Escudero 2003). Higher N concentration in fine roots than coarse roots on July 2 (Table 2) concurred with the results reported by Gordon and Jackson (2000), indicating higher nutrient uptake activity in finer roots. On September 21 no significant differences in N concentration among roots of different diameters (Table 2) may have been due to N retranslocation from fine roots as reported by Nambiar (1987).

Compared to other fertilization treatments on July 2 (Table 3) lower *NRA* and *NRR* with higher *ANDFS* and *RNDFS* of seedlings fertilized with 24 mg·kg<sup>-1</sup> suggested a clear influence of soil N supply on N retranslocation. This contrasts with the con-

clusion that retranslocation is driven by the magnitude of plant nutrient reserves and is unaffected by current supply (Nambiar and Fife 1991; Millard and Proe 1993; Hawkins and Henry 1999; Salifu and Timmer 2001, 2003). Our results, however, concur with van den Driessche (1985) who found an adequate current supply of N reduced the amount of N retranslocation. For Changbai larch seedlings in north-eastern China, limited top-dress applications of N in the first growing season may not facilitate adequate N reserves. Despite the discrepancy among studies associated with different plant materials and methods, we conjecture that sufficient current N supply tends to influence N retranslocation in seedlings with inadequate N reserves.

Negative values of N retranslocation on September 21 in our study (Table 3) contrast with the research results of Salifu and Timmer (2001), where N retranslocation diminished within 120 days after transplanting, but still showed positive values. Our results, however, concur with Imo and Timmer (2001), where current fertilization reduced N retranslocation in all sites and resulted in net assimilation of N (negative values of retranslocation) on low-competition and nutrient poor sites. In early July, soil temperature begins to rise in northeast China, and most seedling roots may have just begun to function effectively for soil nutrient uptake. Then, presumably, N retranslocation was complete before mid September due to low N reserves or because of abundant N content in nursery soil. Larch seedlings may require more N assimilation than retranslocation due to their high growth rates.

Greater biomass and N content in needles and old stems in the 24 kg·ha<sup>-1</sup> N treatment on July 2 (Fig. 1) revealed benefits of this treatment in terms of N accumulation during the growing season. Higher N utilization efficiencies of whole seedlings (Fig. 2A) and CSR (Fig. 2B) and higher fertilizer efficiency (Fig. 2C, D) on September 21 in treatments with 24 kg·ha<sup>-1</sup> N than in other treatments were associated with more biomass and N accumulation in needles and old stems (Fig. 1).

On September 21, lower biomass and N content in the three highest N fertilization treatments indicated an excess of N compared to the control and 24 kg·ha<sup>-1</sup> N treatment (Fig. 3). The local operational N application rate of 81 kg·ha<sup>-1</sup> N combined with the already high initial N content of soils, appeared to exceed the N requirement of Changbai larch seedlings (Fig. 3). High nitrate content in local soils may be due to enhanced nitrification in response to decades of continuous N application using ammonium phosphate as fertilizer. Cruz et al. (2009) found low specific denitrification and high specific nitrogenase activities with low N availability in soils as long as 37 years after annual applications of ammonium phosphate. To improve fertilization efficiency of Changbai larch seedlings, it is recommended: (1) to adopt a lower dose of N application than 81 kg·ha<sup>-1</sup> N, for example, 24 kg·ha<sup>-1</sup> N; (2) to use other fertilizers instead of ammonium phosphate, for example, controlled-release fertilizer; (3) to add nitrification inhibitor into soils; (4) to fertilize using nutrient loading techniques or performing organic amendment.

### Acknowledgements

We thank Changji Jiang and Fusen Liu from Jiangmifeng nursery, Forestry Division of Jilin, China for supplying the materials and facilities for the experiment; Chunsheng Bai and Jinwen Liu for helping with sampling; Qichang Zhang, Chunnan Fan and Jining Zheng from Beihua University, Jilin, China, for their help in seedling transport; Yuwei Liu, Jianfeng Dong, Tianyu Bai and Yuan Chai for their help in chemical analyses. Especially, we appreciate Dr. Barbara J Hawkins for her valuable modification of English language and insightful comments on an earlier version of this manuscript. Also the editorial assistance of Mr. Ruihai Chai and reviewing works of the three anonymous experts are acknowledged.

### References

- Bao SD. 2000. *Soil and agricultural chemistry analysis*. Edition III. Beijing: Chinese Agriculture Press, pp. 25–97. (in Chinese)
- Bown HE, Watt MS, Clinton PW, Mason EG. 2010. Influence of ammonium and nitrate supply on growth, dry matter partitioning, N uptake and photosynthetic capacity of *Pinus radiata* seedlings. *Trees*, **24**(6): 1097–1107.
- Chapin III FS, Kedrowski RA. 1983. Seasonal changes in nitrogen and phosphorus fractions and autumn retranslocation in evergreen and deciduous taiga trees. *Ecology*, **64**(2): 376–391.
- Cruz AF, Hamel C, Hanson K, Selles F, Zentner RP. 2009. Thirty-seven years of soil nitrogen and phosphorus fertility management shapes the structure and function of the soil microbial community in a Brown Chernozem. *Plant and Soil*, **315**: 173–184.
- Dyckmans J, Flessa H. 2001. Influence of tree internal N status on uptake and translocation of C and N in beech: a dual <sup>13</sup>C and <sup>15</sup>N labelling approach. *Tree Physiology*, **21**(6): 395–401.
- Evans JR. 1989. Photosynthesis and nitrogen relationships in leaves of C3 plants. *Oecologia*, **78**(1): 9–19.
- Frak EW, Millard P, Le Roux X, Guillaume S, Wendler R. 2002. Coupling sap flow velocity and amino acid concentrations as an alternative method to <sup>15</sup>N labeling for quantifying nitrogen remobilization by walnut trees. *Plant Physiology*, **130**: 1043–1053.
- Gordon WS, Jackson RB. 2000. Nutrient concentrations in fine roots. *Ecology*, **81**: 275–280.
- Hawkins BJ, Henry G. 1999. Nutrient and bud removal affect biomass and nutrient allocation in Douglas-fir and western red cedar. *Tree Physiology*, **19**: 197–203.
- Helmisaari HS. 1992. Nutrient retranslocation within the foliage of *Pinus sylvestris*. *Tree Physiology*, **10**: 45–58.
- Imo M, Timmer VR. 2001. Growth and nitrogen retranslocation of nutrient loaded *Picea mariana* seedlings planted on boreal mixedwood sites. *Canadian Journal of Forest Research*, **31**: 1357–1366.
- Islam MA, Apostol KG, Jacobs DF, Dumroese RK. 2009. Fall fertilization of *Pinus resinosa* seedlings: nutrient uptake, cold hardiness, and morphological development. *Annals of Forest Science*, **66**(7): 704p1–704p9.
- Kaakinen S, Jolkkonen A, Iivonen S, Vapaavuori E. 2004. Growth, allocation and tissue chemistry of *Picea abies* seedlings affected by nutrient supply during the second growing season. *Tree Physiology*, **24**: 707–719.
- Kagawa A, Sugimoto A, Maximov TC. 2006. Seasonal course of translocation, storage and remobilization of <sup>13</sup>C pulse-labeled photoassimilate in naturally growing *Larix gmelinii* saplings. *New Phytologist*, **171**: 793–804.
- Lehtilä K, Haukioja E, Kaitaniemi P, Laine PA. 2000. Allocation of resources within mountain birch canopy after simulated winter browsing. *Oikos*, **90**: 160–170.
- Lutze JL, Gifford RM. 1998. Acquisition and allocation of carbon and nitrogen by *Danthonia richardsonii* in response to restricted nitrogen supply and CO<sub>2</sub> enrichment. *Plant, Cell & Environment*, **21**: 1133–1141.
- Mead DJ, Preston CM. 1994. Distribution and retranslocation of <sup>15</sup>N in lodgepole pine over eight growing seasons. *Tree Physiology*, **14**: 389–402.
- Miller HG, Cooper JM, Miller JD, Pauline OJ. 1979. Nutrient cycles in pines and their adaptation to poor soils. *Canadian Journal of Forest Research*, **9**: 19–26.
- Millard P, Neilsen, G.H. 1989. The influence of nitrogen supply on the uptake and remobilization of stored N for the seasonal growth of apple trees. *Annals of Botany*, **63**: 301–309.
- Millard P, Proe MF. 1993. Nitrogen uptake, partitioning and internal cycling in *Picea sitchensis* (Bong.) Carr. as influenced by nitrogen supply. *New Phytologist*, **125**: 113–119.
- Millard P, Wendler R, Hepburn A, Smith A. 1998. Variations in the amino acid composition of xylem sap of *Betula pendula* Roth. trees due to remobilization of stored N in the spring. *Plant, Cell & Environment*, **21**: 715–722.
- Millard P, Hester A, Wendler R, Baillie G. 2001. Interspecific defoliation responses of trees depend on sites of winter nitrogen storage. *Functional Ecology*, **15**: 535–543.
- Millard P, Grelet GA. 2010. Nitrogen storage and remobilization by trees: ecophysiological relevance in a changing world. *Tree Physiology*, **30**: 1083–1095.
- Nambiar EKS. 1987. Do nutrients retranslocate from fine roots? *Canadian Journal of Forest Research*, **17**: 913–918.

- Nambiar EKS, Fife DN. 1987. Growth and nutrient retranslocation in needles of radiata pine in relation to nitrogen supply. *Annals of Botany*, **60**: 147–156.
- Nambiar EKS, Fife DN. 1991. Nutrient retranslocation in temperate conifers. *Tree Physiology*, **9**: 185–207.
- Salifu KF, Timmer VR. 2001. Nutrient retranslocation response of *Picea mariana* seedlings to nitrogen supply. *Soil Science and Society of America Journal*, **65**: 905–913.
- Salifu KF, Timmer VR. 2003. Nitrogen retranslocation response of young *Picea mariana* to nitrogen-15 supply. *Soil Science and Society of America Journal*, **67**: 309–318.
- Salifu KF, Apostol KG, Jacobs DF, Islam MA. 2008. Growth, physiology, and nutrient retranslocation in nitrogen-15 fertilized *Quercus rubra* seedlings. *Annals of Forest Science*, **65**(1): 101–101.
- Salifu KF, Islam MA, Jacobs DF. 2009. Retranslocation, plant, and soil recovery of nitrogen-15 applied to bareroot black walnut seedlings. *Communications in Soil Science and Plant Analysis*, **40**: 1408–1417.
- Silla F, Escudero A. 2003. Uptake, demand and internal cycling of nitrogen in saplings of Mediterranean *Quercus* species. *Oecologia*, **136**: 28–36.
- Ueda MU, Mizumachi E, Tokuchi N. 2009. Allocation of nitrogen within the crown during leaf expansion in *Quercus serrata* saplings. *Tree Physiology*, **29**: 913–919.
- van den Driessche R. 1984. Nutrient storage, retranslocation and relationship of stress to nutrition. In: Bowen, G.D., Nambiar, E.K.S. (eds), *Nutrient of Plantation Forests*. New York: Academic Press, pp. 181–209.
- Van den Driessche R. 1985. Late-season fertilization, mineral nutrient reserves, and retranslocation in planted Douglas-fir (*Pseudotsuga menziesii* (Mirb.) Franco) seedlings. *Forest Science*, **31**: 485–496.
- Weinbaum S, Van Kessel, C. 1998. Quantitative estimates of uptake and internal cycling of  $^{14}\text{N}$ -labeled fertilizer in mature walnut trees. *Tree Physiology*, **18**: 795–801.
- Welker JM, Gordon DR, Rice KJ. 1991. Capture and allocation of nitrogen by *Quercus douglasii* seedlings in competition with annual and perennial grasses. *Oecologia*, **87**: 459–466.
- Zhu JJ, Liu ZG, Wang HX, Yan QL, Fang HY, Hu LL, Yu LZ. 2008. Effects of site preparation on emergence and early establishment of *Larix olgensis* in montane regions of northeastern China. *New Forests*, **36**: 247–260.